

Hölder Inequalities and Bounds on the Masses of Light Quarks

T.G. Steele, K. Kostuik, J. Kwan
*Department of Physics and Engineering Physics and
 Saskatchewan Accelerator Laboratory
 University of Saskatchewan
 Saskatoon, Saskatchewan S7N 5C6, Canada.*

February 1, 2008

Abstract

QCD Laplace sum-rules must satisfy a fundamental (Hölder) inequality if they are to consistently represent an integrated hadronic cross-section. After subtraction of the pion-pole, the Laplace sum-rule of pion currents is shown to violate this fundamental inequality unless the u and d quark masses (respectively denoted by m_u and m_d) are sufficiently large, placing a lower bound on the 1GeV $\overline{\text{MS}}$ running masses.

QCD Laplace sum-rules, a technique which equates a theoretical quantity to an integrated hadronic cross-section to determine a QCD prediction of hadronic properties [1], have demonstrated their utility in numerous applications to hadronic physics. It has recently been shown that Laplace sum-rules must satisfy a fundamental inequality in order to consistently represent an integrated hadronic cross-section [2]. When applied to the Laplace sum-rule related to the pion, a strong dependence on the light quark masses m_u and m_d occurs in the analysis of this inequality because the pseudoscalar correlation is proportional to $(m_u + m_d)^2$. In this paper we employ this fundamental Hölder inequality to place lower bounds on the sum of the u and d 1.0 GeV $\overline{\text{MS}}$ running masses.

Consider the correlation function of pseudoscalar currents with the quantum numbers of the pion:

$$J_5(x) = \frac{1}{\sqrt{2}} (m_u + m_d) [\bar{u}(x) i\gamma_5 u(x) - \bar{d}(x) i\gamma_5 d(x)] \quad (1)$$

$$\Pi_5(Q^2) = i \int d^4x e^{iq \cdot x} \langle O | T J_5(x) J_5(0) | O \rangle \quad (2)$$

To leading order in the quark masses, the perturbative contributions Π_5^{pert} are known to four-loop order in the $\overline{\text{MS}}$ scheme [3].

$$\begin{aligned} \Pi_5^{pert}(Q^2) = & \frac{3Q^2}{8\pi^2} (m_u + m_d)^2 \log\left(\frac{Q^2}{\mu^2}\right) \\ & \left[1 + \frac{\alpha}{\pi} (a_0 + a_1 L) + \left(\frac{\alpha}{\pi}\right)^2 [b_0 + b_1 L + b_2 L^2] + \left(\frac{\alpha}{\pi}\right)^3 [c_0 + c_1 L + c_2 L^2 + c_3 L^3] \right] \end{aligned} \quad (3)$$

$$L \equiv \log\left(\frac{Q^2}{\mu^2}\right) \quad , \quad a_0 = \frac{17}{3} \quad , \quad a_1 = -1 \quad (4)$$

$$b_0 = 45.846 \quad , \quad b_1 = -\frac{95}{6} \quad , \quad b_2 = \frac{17}{12} \quad (5)$$

$$c_0 = 465.8463 \quad , \quad c_1 = -194.2393 \quad , \quad c_2 = 38.1667 \quad , \quad c_3 = -2.3020825 \quad (6)$$

Divergent polynomials in Q^2 are ignored in (3) since they correspond to subtraction constants in dispersion relations which do not contribute to sum-rules.

In the QCD sum-rule approach, nonperturbative effects are parametrized by the QCD condensates representing infinite correlation-length vacuum effects [1]. In addition to the QCD condensate contributions, scalar and pseudoscalar correlation functions must also take into account the effects of instantons which represent vacuum effects

with a finite correlation length [4]. Thus the nonperturbative contributions to Π_5 consist of the QCD condensate effects Π_5^{cond} and instanton effects Π_5^{inst} .

To leading order in the quark mass, Π_5^{cond} is given by [5, 6]

$$\Pi_5^{cond}(Q^2) = (m_u + m_d)^2 \left[\frac{\langle \alpha G^2 \rangle}{8\pi Q^2} - \frac{\langle m\bar{q}q \rangle}{Q^2} + \frac{\pi \langle \mathcal{O}_6 \rangle}{4Q^4} \right] \quad (7)$$

where we have used $SU(2)$ symmetry for the dimension-four quark condensates (*i.e.* $(m_u + m_d)\langle \bar{u}u \rangle + \bar{d}d \rangle \equiv 4m\langle \bar{q}q \rangle$), and $\langle \mathcal{O}_6 \rangle$ denotes the dimension six quark condensates

$$\begin{aligned} \langle \mathcal{O}_6 \rangle \equiv & \alpha_s \left[(2\langle \bar{u}\sigma_{\mu\nu}\gamma_5 T^a u \bar{u}\sigma^{\mu\nu}\gamma_5 T^a u \rangle + u \rightarrow d) - 4\langle \bar{u}\sigma_{\mu\nu}\gamma_5 T^a u \bar{d}\sigma^{\mu\nu}\gamma_5 T^a d \rangle \right. \\ & \left. + \frac{2}{3} \langle (\bar{u}\gamma_\mu T^a u + \bar{d}\gamma_\mu T^a d) \sum_{u,d,s} \bar{q}\gamma^\mu T^a q \rangle \right] \end{aligned} \quad (8)$$

The vacuum saturation hypothesis [1] will be used as a reference value for $\langle \mathcal{O}_6 \rangle$

$$\langle \mathcal{O}_6 \rangle = f_{vs} \frac{448}{27} \alpha \langle \bar{q}q\bar{q}q \rangle = f_{vs} 3 \times 10^{-3} \text{GeV}^6 \quad (9)$$

where $f_{vs} = 1$ for exact vacuum saturation. Larger values of effective dimension-six operators found in [7, 8] imply that f_{vs} could be as large as $f_{vs} = 2$. The quark condensate is determined by the GMOR relation [9]

$$(m_u + m_d) \langle \bar{u}u + \bar{d}d \rangle = 4m\langle \bar{q}q \rangle = -2f_\pi^2 m_\pi^2 \quad (10)$$

where $f_\pi = 93 \text{ MeV}$. We will use recent determinations of the gluon condensate $\langle \alpha G^2 \rangle$ [10]

$$\langle \alpha G^2 \rangle = (0.07 \pm 0.01) \text{ GeV}^4 \quad (11)$$

However, it should be noted that there is some discrepancy between [10] and the smaller value $\langle \alpha G^2 \rangle = (0.047 \pm 0.014) \text{ GeV}^4$ found in [8].

The direct single-instanton contribution Π_5^{inst} in the instanton liquid model is [4]

$$\Pi_5^{inst}(Q^2) = \frac{3}{2\pi^2} (m_u + m_d)^2 Q^2 \left[K_{-1} \left(\rho_c \sqrt{Q^2} \right) \right]^2 \quad (12)$$

where $K_n(x)$ denotes the modified Bessel function and $\rho_c \approx 1/(600 \text{ MeV})$ is a fundamental scale in the instanton liquid model [4].

The correlation function $\Pi_5(Q^2)$ satisfies a twice-subtracted dispersion relation

$$\Pi_5(Q^2) = \Pi_5(0) - Q^2 \Pi_5'(0) + \frac{Q^4}{\pi} \int_0^\infty \frac{Im \Pi_5(t)}{t^2(t+Q^2)} dt \quad (13)$$

$$\Pi_5(Q^2) = \Pi_5^{pert}(Q^2) + \Pi_5^{cond}(Q^2) + \Pi_5^{inst}(Q^2) \quad (14)$$

Laplace sum-rules are formed from the dispersion relation by applying the Borel transform operator $\hat{B}$ to (13)

$$\hat{B} = \lim_{\substack{N \rightarrow \infty, \\ Q^2/N \equiv M^2}} \frac{1}{\Gamma(N)} (-Q^2)^N \left(\frac{d}{dQ^2} \right)^N \quad (15)$$

This results in the Laplace sum-rule which exponentially suppresses the high-energy region.

$$\mathcal{R}_0(M^2) \equiv M^2 \hat{B} [\Pi_5(Q^2)] \quad (16)$$

$$\mathcal{R}_0(M^2) = \frac{1}{\pi} \int_0^\infty Im \Pi_5(t) e^{-t/M^2} dt \quad (17)$$

Using the expressions (3, 7, 12, 14) for the correlation function, the definition of $\hat{B}$ and the results of [11], the sum-rule $\mathcal{R}_0(M^2)$ is obtained

$$\begin{aligned}\mathcal{R}_0(M^2) &= \frac{3(m_u + m_d)^2 M^4}{8\pi^2} \left(1 + 4.821098 \frac{\alpha}{\pi} + 21.97646 \left(\frac{\alpha}{\pi} \right)^2 + 53.14179 \left(\frac{\alpha}{\pi} \right)^3 \right) \\ &+ (m_u + m_d)^2 \left(-\langle m\bar{q}q \rangle + \frac{1}{8\pi} \langle \alpha G^2 \rangle + \frac{\pi \langle \mathcal{O}_6 \rangle}{4M^2} \right) \\ &+ (m_u + m_d)^2 \frac{3\rho_c^2 M^6}{8\pi^2} e^{-\rho_c^2 M^2/2} [K_0 (\rho_c^2 M^2/2) + K_1 (\rho_c^2 M^2/2)]\end{aligned}\quad (18)$$

Renormalization group improvement of (18) implies that α , m_u and m_d are running quantities evaluated at the mass scale M in the $\overline{\text{MS}}$ scheme [12].

The running coupling can be directly related to the experimental value [13]

$$\alpha(M_Z) = 0.119 \pm 0.002 \quad (19)$$

by using (19) as an initial condition for the evolution of α to the scale M via the 4-loop $\overline{\text{MS}}$ beta function [14]

$$\mu^2 \frac{d\alpha}{d\mu^2} = \beta \left(\frac{\alpha}{\pi} \right) = -a^2 \sum_{i=0}^{\infty} \beta_i \left(\frac{\alpha}{\pi} \right)^i, \quad a \equiv \frac{\alpha}{\pi} \quad (20)$$

$$\beta_0 = \frac{11 - \frac{2}{3}n_f}{4}, \quad \beta_1 = \frac{102 - \frac{38}{3}n_f}{16}, \quad \beta_2 = \frac{\frac{2857}{2} - \frac{5033}{18}n_f + \frac{325}{54}n_f^2}{64} \quad (21)$$

$$\beta_3 = 114.23033 - 27.133944n_f + 1.5823791n_f^2 - 5.8566958 \times 10^{-3}n_f^3 \quad (22)$$

The only subtlety in this approach is the location of flavour thresholds where the number of effective flavour degrees of freedom n_f change. In general, matching conditions must be imposed at these thresholds to relate QCD with n_f quarks to an effective theory with $n_f - 1$ light quarks and a decoupled heavy quark [15]. Using the matching threshold μ_{th} defined by $m_q(\mu_{th}) = \mu_{th}$, where m_q is the running quark mass, the matching condition to three-loop order is [16]

$$a^{(n_f-1)}(\mu_{th}) = a^{(n_f)}(\mu_{th}) \left[1 + 0.1528 \left[a^{(n_f)}(\mu_{th}) \right]^2 + \{0.9721 - 0.0847(n_f - 1)\} \left[a^{(n_f)}(\mu_{th}) \right]^3 \right] \quad (23)$$

For energies above the charm threshold, $\alpha(M)$ depends only on the initial condition (19) and the bottom threshold. The currently accepted range for the bottom threshold [13]

$$4.1 \text{ GeV} \leq m_b(m_b) \leq 4.4 \text{ GeV} \quad (24)$$

has a negligible effect on the evolution in $\alpha(M)$ in comparison with the uncertainty (19) in $\alpha(M_Z)$.

Similarly, the running quark masses can be parametrized by their (unknown) value at 1 GeV. Defining

$$m = \frac{1}{2} (m_u + m_d) \quad (25)$$

$$w(M) = \frac{m(M)}{m(1 \text{ GeV})}, \quad w(1 \text{ GeV}) = 1 \quad (26)$$

allows the quark mass at the scale M to be determined via the four-loop $\overline{\text{MS}}$ scheme anomalous mass dimension [17]

$$\frac{dw}{d\mu} = w(\mu) \frac{\gamma \left[\frac{\alpha(\mu)}{\pi} \right]}{\beta \left[\frac{\alpha(\mu)}{\pi} \right]} \quad (27)$$

$$\gamma(x) = -x[1 + \sum_{i=1} \gamma_i x^i] \quad (28)$$

$$\gamma_1 = 4.20833 - 0.138889n_f \quad , \quad \gamma_2 = 19.5156 - 2.28412n_f - 0.0270062n_f^2 \quad (29)$$

$$\gamma_3 = 98.9434 - 19.1075n_f + 0.276163n_f^2 - 0.00579322n_f^3 \quad (30)$$

To evaluate the quark mass above the charm threshold the currently accepted values in the $\overline{\text{MS}}$ scheme [13] will be used

$$1.1 \text{ GeV} \leq m_c(m_c) \leq 1.4 \text{ GeV} \quad (31)$$

In sum-rule applications, the M dependence of the theoretical result $\mathcal{R}_0(M^2)$ (18) over a range of M values is used to predict the resonance parameters contained in $Im\Pi(t)$ through the relation (17). Since $Im\Pi(t)$ is in general related to a physical cross-section, fundamental inequalities for Laplace sum-rules can be constructed using the property that $Im\Pi(t) > 0$. Thus the simplest inequality that can be obtained is

$$\mathcal{R}_0(M^2) = \int Im\Pi(t) e^{-t/M^2} dt \geq 0 \longrightarrow \mathcal{R}_0(M^2) \geq 0 \quad (32)$$

More stringent sum-rule inequalities can be developed using integral inequalities. In particular, Hölder inequalities for Laplace sum-rules have been shown to be valuable in studying self-consistency of QCD sum-rules and in obtaining bounds on the electromagnetic polarizability of charged pions [2]. Hölder's inequality for integrals defined over a measure $d\mu$ is [18]

$$\left| \int_{t_1}^{t_2} f(t)g(t)d\mu \right| \leq \left(\int_{t_1}^{t_2} |f(t)|^p d\mu \right)^{1/p} \left(\int_{t_1}^{t_2} |g(t)|^q d\mu \right)^{1/q}, \quad (33)$$

$$\frac{1}{p} + \frac{1}{q} = 1; \quad p, q \geq 1.$$

When $p = q = 2$ the Hölder inequality reduces to the well known Schwarz inequality. The key idea in applying Hölder's inequality to sum-rules is recognizing that since $Im\Pi(t)$ is positive it can serve as the measure $d\mu = Im\Pi(t)dt$ in (33). With the definitions

$$\tau = \frac{1}{M^2} \quad (34)$$

$$\mathcal{S}_k(\tau) \equiv \int_{\mu_{th}}^{\infty} Im\Pi(t) t^k e^{-t\tau} dt \quad (35)$$

(33) with $d\mu = Im\Pi(t)dt$, $f(t) = t^\alpha e^{-at\tau}$, $g(t) = t^\beta e^{-bt\tau}$, leads to the following inequalities for $\mathcal{S}_k(\tau)$

$$\mathcal{S}_{\alpha+\beta}(\tau) \leq \mathcal{S}_{\alpha p}^{1/p}(ap\tau) \mathcal{S}_{\beta q}^{1/q}(bq\tau); \quad a + b = 1. \quad (36)$$

Imposing restrictions that we have the integer value $k = 0$ in our sum-rule (17) leads to the following inequality.

$$\mathcal{S}_0[\omega\tau_{min} + (1-w)\tau_{max}, s_0] \leq \mathcal{S}_0^\omega[\tau_{min}, s_0] \mathcal{S}_0^{1-\omega}[\tau_{max}, s_0], \quad (37)$$

$$0 \leq \omega \leq 1; \quad \delta\tau = \tau_{max} - \tau_{min} > 0 \quad (38)$$

If the pion pole contribution to $Im\Pi_5(t)$ is explicitly included, then the sum-rule (17) becomes

$$\mathcal{R}_0(\tau) = 2f_\pi^2 m_\pi^4 e^{-m_\pi^2 \tau} + \int_{m_\pi^2}^{\infty} Im\Pi_5(t) e^{-t\tau} dt \quad (39)$$

where m_π represents the pion mass and $f_\pi = 93 \text{ MeV}$ is the pion decay constant. Since for any reasonable range of τ , $\exp(-m_\pi^2 \tau) \approx 1$, we find

$$\mathcal{S}_0(\tau) \equiv \mathcal{R}_0(\tau) - 2f_\pi^2 m_\pi^4 = \int_{m_\pi^2}^{\infty} \text{Im}\Pi_5(t) e^{-t\tau} dt \quad (40)$$

Thus if the sum-rule is a valid and consistent representation of the integration of $\text{Im}\Pi(t)$ in (17) then after subtraction of the pion pole, the sum-rule $\mathcal{S}_0(\tau)$ must satisfy the fundamental inequality (37).

$$\rho_0 \equiv \frac{\mathcal{S}_0[\tau + (1-\omega)\delta\tau]}{\mathcal{S}_0^\omega[\tau]\mathcal{S}_0^{1-\omega}[\tau + \delta\tau]} \leq 1 \quad , \quad \forall 0 \leq \omega \leq 1 \quad (41)$$

Provided that $\delta\tau$ is reasonably small (in QCD $\delta\tau \approx 0.1 \text{ GeV}^{-2}$ appears sufficient) these inequalities are insensitive to the value of $\delta\tau$, permitting a simple analysis of the inequality as a function of the energy-scale M .

The first applications of inequalities to quark-mass bounds used the simple positivity constraint $\mathcal{S}_0(\tau) \geq 0$ [5, 12, 19]. We extend this analysis by using the more stringent Hölder inequality, higher-loop perturbative corrections, and inclusion of instanton effects which are important for the pseudoscalar channel.

For a fixed value of $M = 1/\sqrt{\tau}$ the Hölder inequality can be used to find the minimum value of $m(1 \text{ GeV}) \equiv [m_u(1 \text{ GeV}) + m_d(1 \text{ GeV})]/2$ for which the inequality (41) is satisfied. This lower bound on $m(1 \text{ GeV})$ depends on the energy scale M as indicated in Figure 1. However, the increasingly large coefficients in the perturbative portion of (18) suggests that these mass bounds could depend strongly upon higher-order perturbative effects. Asymptotic Padé approximation methods [20] applied to the perturbative part of (18) result in the following estimate of $\mathcal{R}_0(M^2)$ with perturbative effects to five-loop order.

$$\begin{aligned} \mathcal{R}_0(M^2) = & \frac{3(m_u + m_d^2)^2 M^4}{8\pi^2} \left(1 + 4.821098 \frac{\alpha}{\pi} + 21.97646 \left(\frac{\alpha}{\pi} \right)^2 + 53.14179 \left(\frac{\alpha}{\pi} \right)^3 + 137.6 \left(\frac{\alpha}{\pi} \right)^4 \right) \\ & + (m_u + m_d)^2 \left(-\langle m\bar{q}q \rangle + \frac{1}{8\pi} \langle \alpha G^2 \rangle + \frac{\pi \langle \mathcal{O}_6 \rangle}{4M^2} \right) \\ & + (m_u + m_d)^2 \frac{3\rho_c^2 M^6}{8\pi^2} e^{-\rho_c^2 M^2/2} [K_0 (\rho_c^2 M^2/2) + K_1 (\rho_c^2 M^2/2)] \end{aligned} \quad (42)$$

By comparing mass bounds with and without inclusion of this estimated five-loop perturbative contribution, the uncertainty in the mass bounds of Figure 1 devolving from truncation of the perturbative expansion can be estimated. In addition to this perturbative uncertainty, a 50% uncertainty in the vacuum saturation hypothesis (parameterized by f_{vs}), a 15% uncertainty in the instanton size ρ_c , and the effects of varying the input parameters within the ranges (11,19,24,31) must also be considered. Figure 2 illustrates the effect of all sources of uncertainty upon the mass bounds obtained from the Hölder inequality. As indicated by the figure, the analysis is extremely stable.

The increase of the quark mass bound m_{min} with decreasing M implies that the minimum energy scale M at which the pseudoscalar sum-rule is considered valid will provide the most stringent bound on the quark mass m . The analysis of the τ hadronic width [21], and hadronic contributions to $\alpha_{EM}(M_Z)$ and the anomalous magnetic moment of the muon [22] provides excellent evidence for the validity of QCD sum-rule methods at the energy scale $M_\tau \approx 1.8 \text{ GeV}$. Furthermore, the largest value of M at which the uncertainties in the analysis are visible in Figure 2 corresponds to $M \approx M_\tau$, providing support for the validity of the mass bounds at $M = M_\tau$. Thus a valid and *conservative* bound on the 1.0 GeV $\overline{\text{MS}}$ quark masses is obtained from Figure 2 at $M = M_\tau$

$$m(1 \text{ GeV}) = \frac{1}{2} [m_u(1 \text{ GeV}) + m_d(1 \text{ GeV})] \geq 3 \text{ MeV} \quad (43)$$

This conservative bound is phenomenologically significant since the Particle Data Group quotes a lower bound of $m(1 \text{ GeV}) \geq 2.25 \text{ MeV}$ [13].

Acknowledgements: TGS is grateful for the financial support of the Natural Sciences and Engineering Research Council of Canada (NSERC).

References

- [1] M.A. Shifman, A.I. Vainshtein, V.I. Zakharov, Nucl. Phys. **B147** (1979) 385.
- [2] M. Benmerrouche, G. Orlandini, T.G. Steele, Phys. Lett. **B356** 573, (1995);
M. Benmerrouche, G. Orlandini, T.G. Steele, Phys. Lett. **B366** 354, (1996);
T.G. Steele, S. Alavian, J. Kwan, Phys. Lett. **B392** (1997) 189.
- [3] K.G. Chetyrkin, Phys. Lett. **B390** (1997) 309.
- [4] E.V. Shuryak, Nucl. Phys. **B214** (1983) 237.
- [5] C. Becchi, S. Narison, E. de Rafael, F.J. Yndurain, Z. Phys. **C8** (1981) 335.
- [6] E. Bagán, J.I. Latorre, P. Pascual, Z. Phys. **C32**,43 (1986).
- [7] C.A. Dominguez, J. Sola, Z. Phys. **C40** (1988) 63.
- [8] V. Gimenez, J. Bordes, J.A. Penarrocha, Nucl. Phys. **B357** (1991) 3.
- [9] M. Gell-Mann, R. J. Oakes and B. Renner, Phys. Rev. **2195** (1968) 175.
- [10] S. Narison, Nucl. Phys. Proc. Suppl. **54A** (1997) 238.
- [11] A.E. Dorokhov, S.V. Esaibegian, N.I. Kochelev and N.G. Stefanis, J. Phys. **G23** (1997) 643.
- [12] S. Narison, E. de Rafael, Phys. Lett. **B103** (1981) 57.
- [13] Particle data Group, C. Caso *et al*, Eur. Phys. J. **C3** (1998) 1.
- [14] T. van Ritbergen, J.A.M. Vermaseren, S.A. Larin, Phys. Lett. **B400** (1997) 379
- [15] G. Rodrigo, A. Santamaria, Phys. Lett. **B313** (1993) 441.
- [16] K.G. Chetyrkin, B.A. Kniehl, M. Steinhauser, Nucl. Phys. **B510** (1998) 61.
- [17] K.G. Chetyrkin, Phys. Lett. **B404** (1997) 161.
- [18] E.F. Beckenbach, R. Bellman, *Inequalities*, (Springer, Berlin, 1961);
S.K. Berberian, *Measure and Integration*, (MacMillan, New York, 1965).
- [19] L. Lellouch, E. de Rafael, J. Taron, Phys. Lett. **B414** (1997) 195-204.
- [20] J. Ellis, I. Jack, D.R.T. Jones, M. Karliner, M.A. Samuel, Phys. Rev. **D57** (1998) 2665;
M.A. Samuel, J. Ellis, M. Karliner, Phys. Lett. **B400** (1997) 176;
V. Elias, T.G. Steele, F. Chishtie, R. Migneron, K. Sprague, Phys. Rev. **D58** (1998) 116007.
- [21] E. Braaten, S. Narison, A. Pich, Nucl. Phys. **B373** (1992) 581.
- [22] M. Davier, A. Höcker, Phys. Lett. **B435** (1998) 427.




Figure 1: The quantity $m_{\min}$, representing the minimum value of m for which the Hölder inequality (41) is satisfied, is plotted as a function of the mass scale M using central values of all input parameters.




Figure 2: The effect of uncertainties in the input parameters on $m_{\min}$ is plotted as a function of the mass scale M . Higher-loop effects are estimated by the inclusion of the five-loop perturbative correction in (42).